

# A One-Page Summary Report of Genome Sequencing for the Healthy Adult

Jason L. Vassy<sup>a, d, e</sup> Heather L. McLaughlin<sup>f, g</sup> Calum A. MacRae<sup>b-d</sup>  
Christine E. Seidman<sup>b-d</sup> Denise Lautenbach<sup>b</sup> Joel B. Krier<sup>b, d</sup> William J. Lane<sup>f</sup>  
Isaac S. Kohane<sup>d</sup> Michael F. Murray<sup>h</sup> Amy L. McGuire<sup>i</sup> Heidi L. Rehm<sup>f, g</sup>  
Robert C. Green<sup>b, d, g</sup> for the MedSeq Project



Divisions of <sup>a</sup>General Internal Medicine and Primary Care, <sup>b</sup>Genetics and <sup>c</sup>Cardiology, Department of Medicine, Brigham and Women's Hospital, <sup>d</sup>Harvard Medical School, <sup>e</sup>VA Boston Healthcare System, <sup>f</sup>Department of Pathology, Brigham and Women's Hospital and Harvard Medical School, and <sup>g</sup>Partners Personalized Medicine, Boston, Mass., <sup>h</sup>Genomic Medicine Institute, Geisinger Health System, Danville, Pa., and <sup>i</sup>Center for Medical Ethics and Health Policy, Baylor College of Medicine, Houston, Tex., USA

## Key Words

Whole-genome sequencing · Genomic screening · Laboratory reporting · Molecular testing

## Abstract

As genome sequencing technologies increasingly enter medical practice, genetics laboratories must communicate sequencing results effectively to nongeneticist physicians. We describe the design and delivery of a clinical genome sequencing report, including a one-page summary suitable for interpretation by primary care physicians. To illustrate our preliminary experience with this report, we summarize the genomic findings from 10 healthy participants in a study of genome sequencing in primary care. © 2015 S. Karger AG, Basel

## Introduction

The clinical implementation of genomic medicine has begun and is widely anticipated to bring exciting medical benefits. Whole-exome and whole-genome sequencing, hereafter referred to as genomic sequencing (GS), is be-

coming increasingly useful in the diagnosis and treatment of rare diseases [1–3] and cancer [4–6] and is currently used predominantly by genetics specialists or subspecialty practitioners [1]. In anticipation of the more widespread introduction of clinical sequencing to general medicine practice, we are conducting a randomized clinical trial studying the impact of GS, with attendant incidental findings, in both the specialty care of a hereditary illness and the primary care of healthy adults. In order to study the impact of GS and its clinical outcomes in this context, we have refined an interpretive pipeline for the filtration and analysis of any medically relevant variants discovered in an individual genome [7] and designed a report to communicate these results effectively to nongeneticist physicians. Here, we describe this report and our early experiences delivering GS results to primary care physician participants via a Genome Report (fig. 1; online suppl. Appendix, see [www.karger.com/doi/10.1159/000370102](http://www.karger.com/doi/10.1159/000370102)), along with the genomic results from the first 10 healthy participants.

H.L.R. and R.C.G. contributed equally to this work.

## The Challenge of Genomic Screening in Healthy Individuals

GS of individuals without a suspected genetic illness is currently not recommended [8] but may eventually be utilized in population-based genomic screening for the practice of preventive medicine [9, 10]. Under this model, knowledge gained from GS might guide heightened surveillance in the face of increased genetic predisposition, reveal pharmacogenomic information that could maximize drug efficacy and minimize adverse effects, or be used for reproductive planning. As one of the NIH-funded centers in the Clinical Sequencing Exploratory Research Consortium [11], we are conducting the MedSeq Project to develop such processes and to explore the risks and benefits of sequencing healthy adults [12]. A significant challenge for genomic screening with GS is to identify, distill, and report the variants that have clinical significance for a healthy individual.

Molecular laboratories have developed standards and ontologies for analyzing and communicating the clinical significance of genetic variants found during GS of individuals with a suspected genetic disease [13, 14]. After automated filtration and manual curation, geneticists reporting on GS may convey a variant's clinical significance by using a variant classification schema such as 'benign', 'likely benign', 'variant of uncertain significance (VUS)', 'likely pathogenic', or 'pathogenic'. Because many variants fall into the uncertain category, some laboratories will further subclassify VUS into subcategories such as 'favor benign' or 'favor pathogenic'. While efforts for standardization are under way, there is as yet limited consensus on the validity and evidence for pathogenicity that a variant should have before being reported to physicians and patients. If a laboratory identifies a VUS in a gene related to the reason for clinical sequencing, it will likely choose to report that finding to the ordering physician, despite the inherent uncertainties. For example, a VUS in the cardiac  $\beta$ -myosin heavy chain (*MYH7*) gene will likely be reported when the patient has familial hypertrophic cardiomyopathy.

However, when sequencing is performed in healthy individuals without a specific reason for genomic testing, the scenario more closely resembles screening tests used in preventive medicine, which are generally held to higher standards of validity and utility. For example, one might argue that a VUS in the *MYH7* gene should be reported to the patient with familial cardiomyopathy, because additional testing might determine whether the variant segregates with disease in the family. By contrast, whether this variant should be reported to the generally healthy individual without a personal or familial history of cardiomy-

opathy is a much more difficult question, absent studies with long-term follow-up that can help elucidate the pathogenicity of the variant as well as the penetrance of pathogenic variants in an unselected population.

## Interpreting GS Results of Healthy Individuals


As part of the MedSeq Project, we are performing GS in patients who have been identified by their primary care physicians as having no major medical conditions. We are communicating the results of GS to nongeneticist physicians recruited from the primary care practices at our academic medical center, despite the challenges of interpreting and reporting identified variants and the concerns that the primary care workforce may not be prepared for the clinical integration of genomic information [15–18]. For each participant, whole-genome sequencing is performed in a Clinical Laboratory Improvement Amendments (CLIA)-approved laboratory and the resulting raw files of annotated high-quality variants are filtered based on their allele frequencies, previous disease associations in reference databases, genic locations, and predicted functional impacts. Variants are then manually classified by a review of genetic and functional evidence from the scientific literature as previously described [7]. At the end of this pipeline, the Genome Report for generally healthy individuals includes variants classified as 'likely pathogenic', 'pathogenic', or VUS subclassified as 'favor pathogenic'. In cases of uncertainty in how best to weight evidence for variant classification, the larger study team discusses the evidence as a group, careful to adhere to this systematic approach for variant classification and returning results. In this study, the laboratory interprets the GS data and issues the Genome Reports without any clinical knowledge about the patient. Given the low pretest probability of disease among healthy individuals, we do not report the vast majority of VUSs and none of the 'benign' or 'likely benign' variants. By reporting variants with at least some degree of evidence favoring pathogenicity, we aim to strike a balance between the risks of returning uncertain results and the potential clinical benefits for the generally healthy person undergoing genomic screening.

## Communicating GS Results to Physicians


The molecular and clinical geneticists, genetic counselors, bioinformaticians, clinicians, and social scientists comprising the study team have been guided by two main

**Monogenic disease risk:** This section includes rare variants classified as *pathogenic*, *likely pathogenic*, or, occasionally, *uncertain significance*: favor *pathogenic* in any of several thousand Mendelian disease genes where risk of disease is predicted.

**LABORATORY FOR MOLECULAR MEDICINE**  
95 Landsdowne St, Cambridge, MA 02139  
Phone: (617) 768-8500 / Fax: (617) 768-8513  
<http://pcpgm.partners.org/lmm>



**CENTER FOR PERSONALIZED GENETIC MEDICINE**

A teaching affiliate of  
 HARVARD MEDICAL SCHOOL

**Name: DOE, JONATHAN**      Accession ID: **PMXX-12345**  
**DOB: 12/34/5678**      MRN: **123456780**      Family #: **F12345**  
**Sex: Male**      Specimen: **Blood, Peripheral**      Referring physician: **MedSeq**  
**Race: Caucasian**      Received: **05/03/2013**      Referring facility: **MedSeq**  
**Indication for testing: MedSeq, Primary Care**      Test: **WGS-pnIA, SeqConV2, WGS-GGR**

### GENOME REPORT

**RESULT SUMMARY**  
Sequencing of this individual's genome was performed and covered 95.7% of all positions at 8X coverage or higher, resulting in over 5.2 million variants compared to a reference genome. These data were analyzed to identify previously reported variants of potential clinical relevance as well as novel variants that could reasonably be assumed to cause disease (see methodology below). All results are summarized on page 1 with further details on subsequent pages.

**MONOGENIC DISEASE RISK: 1 VARIANT IDENTIFIED**  
This test identified 1 genetic variant that may be responsible for existing disease or the development of disease in this individual's lifetime.

Disease Inheritance	Gene Transcript	Zygoty Variant	Classification
Chondrodysplasia punctata X-linked	ARSE NM_000047.2	Hemizygous c.410G>C p.Gly137Ala	Uncertain Significance: Favor Pathogenic

**CARRIER STATUS: 2 VARIANTS IDENTIFIED**  
This test identified carrier status for 2 autosomal recessive disorders.

Disease Inheritance	Gene Transcript	Zygoty Variant	Classification	Carrier Phenotype*
Cystic Fibrosis Autosomal Recessive	CFTR NM_000492.3	Heterozygous c.3846G>A p.Trp1282X	Pathogenic	None reported
Glycogen storage disease 7 Autosomal recessive	PFKM NM_000289.5	Heterozygous c.2371G>A	Pathogenic	None reported

As a carrier for recessive genetic variants, this individual is at higher risk for having a child with one or more of these highly penetrant disorders. To determine the risk for this individual's future children to be affected, the partner of this individual would also need to be tested for variants in these genes. Other biologically related family members may also be carriers of these variants. \*Carriers for some recessive disorders may be at risk for certain phenotypes. Please see variant descriptions for more information.

**PHARMACOGENOMIC ASSOCIATIONS**  
This test identified the following pharmacogenomic associations. Additional pharmacogenomic results may be requested, but will require additional molecular confirmation prior to disclosure.

Drug	Risk and Dosing Information
Warfarin	Increased dose requirement
Clopidogrel	Typical response to clopidogrel
Digoxin	Intermediate metabolism and serum concentration of digoxin
Metformin	Decreased glycemic response to metformin
Simvastatin	Typical risk of simvastatin-related myopathy

**RED BLOOD CELL AND PLATELET ANTIGENS**  
This test identified the ABO Rh blood type as AB Negative. Based on their results, this person is a very desirable universally compatible platelet donor. Additional RBC and platelet antigen information is available at the end of the report.

It should be noted that the disease risk section of this report is limited only to variants with strong evidence for causing highly penetrant disease, or contributing to highly penetrant disease in a recessive manner. Not all variants identified have been analyzed, and not all regions of the genome have been adequately sequenced. These results should be interpreted in the context of the patient's medical evaluation, family history, and racial/ethnic background. Please note that variant classification and/or interpretation may change over time if more information becomes available. For questions about this report, please contact the Genome Resource Center at [GRC@partners.org](mailto:GRC@partners.org).

**Carrier status:** This section includes heterozygous variants classified as *pathogenic*, *likely pathogenic*, or, occasionally, *uncertain significance*: favor *pathogenic* in genes associated with recessive Mendelian diseases. The presence of only one copy of a pathogenic variant at these loci makes the individual a carrier for the disease, with a risk of having a child with the disease if the other parent is also a carrier. Carrier status for some conditions also have milder phenotypic implications for the carrier himself.

**Pharmacogenomics:** This section includes pharmacogenomic associations for safety and efficacy of the 5 common medications meeting Pharmacogenomics Knowledge Base (PharmGKB) Clinical Annotation Levels of Evidence Classes I & II.

**Blood groups:** This section includes a summary of the genomic predictions of ABO and Rh blood types, along with implications for blood donation and transfusion. Any additional rare red blood cell or platelet antigens are also reported.

**Fig. 1.** Summary page of the Genome Report for GS results for a generally healthy individual.

principles in designing our Genome Report for healthy individuals. First, although genomic information has been heralded as a disruptive technology, we envision that its successful integration into medical practice will conform to standard clinical workflows: through providers to patients. The molecular laboratory sends the Genome Report, via the electronic health record, to the ordering physician. The report is written in medical terminology and not at a lay-person level, thus representing a different paradigm than direct-to-consumer genomics products. The second guiding principle is that GS reports must relay the most important findings concisely to busy physicians, but brevity must not mask complexity when it is present. The reports must be complete while still conveying a sense of prioritization, so that overwhelming

amounts of test information and clinically uncertain results do not distract from the results in each report that may be most clinically significant for the patient. These guiding principles allow us to incorporate feedback flexibly to improve the Genome Report. For example, we eliminated the word 'Mendelian' from an earlier version of the report after nongeneticist physicians reported uncertainty in its meaning.

The first page of the Genome Report (fig. 1) is a summary of the key findings from the GS, while the subsequent pages give more detailed information. This format mirrors the precedent of prioritized information seen in other clinical reports. For example, a radiologist's full text of a chest X-ray report may describe in detail the radiographic appearance of the lungs, cardiac silhouette, ribs,

vertebrae, and overlying soft tissue. However, the ‘Impression’ section of the report, often set apart as separate text with bold lettering, might simply state ‘No acute cardiopulmonary process’ or ‘Right middle lobe pneumonia’. The receiving physician often looks here first and then uses clinical judgment to determine whether the details in the full text of the report warrant further consideration. In our Genome Report, the first page is the genomic ‘Impression’ section, summarizing the identified variants that may cause monogenic disease or that indicate carrier status of autosomal recessive disorders. It also includes brief summaries of the implications of the patient’s genome for blood donation and transfusion [19] and for the safety and efficacy of five medications commonly used in primary care meeting the Pharmacogenomics Knowledge Base Clinical Annotation Levels of Evidence Classes I and II criteria [20]: warfarin, clopidogrel, digoxin, metformin, and simvastatin. The subsequent pages of the Genome Report go on to define the precise variants identified and provide results and interpretation in greater detail (online suppl. Appendix), including a discussion of the scientific evidence favoring or disfavoring pathogenicity for each variant, references to relevant studies, information about the associated diseases, and risk to family members. This section also gives greater detail about the pharmacogenomic and blood antigen loci interrogated in each genome as well as the details and limitations of the sequencing and interpretation methodologies. Despite concerns that non-geneticist physicians are unprepared for genomic medicine [15–18], pilot testing among primary care physician participants of the MedSeq Project suggests that the Genome Report communicates clinical sequencing results effectively. Many physicians expressed the idea that the information is initially daunting but ultimately manageable, represented by one who stated: ‘Although at first glance it is intimidating, I think if you get used to it, you probably can handle this.’

Although we have designed the Genome Report with clarity in mind, we do not assume that it alone is sufficient to turn primary care physicians into proficient genomic medicine practitioners. Primary care physician participants of the MedSeq Project first underwent an orientation to genomics concepts, including inheritance patterns and pharmacogenomics, and to the format of the Genome Report. Comprised of two 1-hour in-person group classes and 4 h of self-paced online modules, these sessions are comparable in length to other continuing medical education opportunities that are a part of routine physician recertification. We also provide individualized support to physicians during the course of the study by offering the

option to contact study-affiliated medical geneticists and genetic counselors for assistance in interpreting and managing their patients’ GS results. We think this model of physician training and support will be scalable for an era when the genomics workforce will be insufficient to meet the clinical demand [21], and, as part of our research, we are collecting data on the use of these resources.

### Sequences from the First 10 Healthy Patients

Table 1 shows the GS results reported for the first 10 primary care patient participants. A total of 211 unique variants were assessed and classified. After assessment, 70 (33%) of these were classified as benign or likely benign, 116 (55%) as VUS, and 25 (12%) as likely pathogenic or pathogenic. We have reported 3 variants associated with monogenic disease risk in 3 healthy participants. The first of these was a variant in the arylsulfatase E (*ARSE*) gene (fig. 1; table 1) previously observed in 2 males with chondrodysplasia punctata, an X-linked disorder of bone and cartilage development. Additionally, prior research indicated that the variant deleteriously impacted protein function. The same variant was identified in a male family member without the disease, though penetrance is known to be incomplete for the disorder. Given this limited scientific evidence, we classified and reported the *ARSE* variant as a ‘VUS, favor pathogenic’. For a generally healthy adult, the clinical significance of this variant is unknown, but by virtue of its appearance on the Genome Report, the primary care clinician has the opportunity to review the patient’s history and physical examination and to ask more directed questions about other family members.

Most variants reported in our healthy patients are associated with a carrier state for an autosomal recessive condition (table 1). These carrier states may be associated with milder phenotypic implications for the carriers themselves, which we include in the Genome Report. For example, in 1 individual we reported a pathogenic variant in the *WFS1* gene associated with Wolfram syndrome, a recessive neurodegenerative disease characterized by sensorineural deafness, type 1 diabetes mellitus, and pituitary gland dysfunction. However, carriers of *WFS1* variants may themselves demonstrate some degree of low-frequency hearing loss and impaired glucose metabolism. Although most patients have had at least 1 carrier variant, we are finding that primary care physicians and their healthy patients understand the extremely low likelihood that these carrier states will impact the health of the patients and their family members.

**Table 1.** GS results reported for the first 10 generally healthy participants in the MedSeq Project

Section of the Genome Report	Results	Classification	
<b>Monogenic disease risk</b> (3 variants identified in 3 patients)	Gene <i>LHX4</i>	Disease Combined pituitary hormone deficiency	Pathogenic
	<i>KCNQ1</i>	Romano-Ward syndrome (a long QT syndrome)	Likely pathogenic
	<i>ARSE</i>	Chondrodysplasia punctata	VUS: favor pathogenic
<b>Carrier status</b> (22 variants identified in 10 patients)	Gene <i>MMACHC</i>	Disease Methylmalonic aciduria and homocystinuria, cblC type	Pathogenic
	<i>CFTR</i>	Cystic fibrosis	Pathogenic
	<i>PFKM</i>	Glycogen storage disease 7	Pathogenic
	<i>CUBN</i>	Imerslund-Gräsbeck syndrome	Pathogenic
	<i>DUOX2</i>	Hypothyroidism	Pathogenic
	<i>ABCA4</i>	Stargardt disease	Pathogenic
	<i>MPO</i>	Myeloperoxidase deficiency	Pathogenic
	<i>BTB</i>	Biotinidase deficiency	Pathogenic
	<i>PYGL</i>	Glycogen storage disease 6	Pathogenic
	<i>SPG7</i>	Spastic paraplegia type 7	Pathogenic
	<i>WFS1</i>	Wolfram syndrome	Pathogenic
	<i>CLRN1</i>	Usher syndrome type III	Pathogenic
	<i>CYP1B1</i>	Primary congenital glaucoma	Pathogenic
	<i>NLRP7</i>	Recurrent hydatidiform mole	Pathogenic
	<i>SPATA7</i>	Leber congenital amaurosis	Likely pathogenic
	<i>ERCC5</i>	Xeroderma pigmentosum	Likely pathogenic
	<i>COL7A1</i>	Epidermolysis bullosa dystrophica	Likely pathogenic
	<i>KCNQ1</i>	Jervell and Lange-Nielsen syndrome	Likely pathogenic
	<i>NAGA</i>	Alpha-N-acetylgalactosaminidase deficiency	Likely pathogenic
	<i>SP110</i>	Hepatic veno-occlusive disease with immunodeficiency	Likely pathogenic
<i>RAB27A</i>	Familial hemophagocytic lymphohistiocytosis	VUS: favor pathogenic	
<i>CNGA3</i>	Achromatopsia	VUS: favor pathogenic	
<b>Pharmacogenomics</b>			
Warfarin	2 patients with <i>decreased</i> predicted dose requirement 3 patients with <i>increased</i> predicted dose requirement		
Clopidogrel	2 patients with <i>increased</i> predicted anti-platelet response		
Digoxin	4 patients with <i>increased</i> predicted serum drug concentration 1 patient with <i>decreased</i> predicted serum drug concentration		
Metformin	5 patients with <i>decreased</i> predicted glycemic response		
Simvastatin	2 patients with <i>increased</i> risk of medication-related myopathy		
<b>Blood groups</b>	3 patients predicted to be <i>desirable platelet donors</i>		

### The Future of the Genome Report

Our early experience in piloting the Genome Report has allowed us to optimize its format and content, including descriptive section header names understandable to primary care physicians and the appropriate distribution of details on the first versus the subsequent pages. More work remains, however. Communication today is instantaneous, and information is updated in real-time. Medical records, however, have generally not kept pace with

state-of-the-art standards for communicating, displaying, and accessing information. At the same time, clinical data are becoming more complex, and the communication of results will need to become more dynamic and prioritized, allowing the physician and patient to access, digest, and use the data to variable degrees over time. One can imagine GS reports that have more creative dashboards for the clinician, and that incorporate hyperlinks to greater detail or to external references, allowing a physician to narrow or expand the focus of interrogating a

patient's genome. With the discovery of new disease genes, mutations, and clinically meaningful gene-gene and gene-exposure interactions, the genome report will need to reflect that new dynamism of genomic data. While our Genome Report provides an accessible summary of complex results using current clinical laboratory reporting and medical practices, it must eventually be adaptable to accurately and concisely communicate the full medical content contained within the genome. We are currently producing the Genome Report using the GeneInsight software suite [22], which is linked to a genomic knowledgebase, so that variant classifications remain up-to-date based on new scientific discovery. Through this process, the physician receives an alert message when a previously reported variant in a patient's genome has been reclassified on the basis of new knowledge (e.g. from 'VUS' to 'likely pathogenic'). One can also imagine a dynamic genome report that evolves to maintain relevance to the patient's individual clinical context over their life course, prioritizing carrier status during reproductive years and highlighting other genomic variants when newly abnormal clinical tests, physical findings, and medications are recorded. An active area of research

is the role patients' preferences might play in shaping which GS results are reported and how. Regardless of the format that GS results take moving forward, physicians will need clear signposts to help them navigate their patients' genomes.

## Acknowledgements

We thank the physician participants and patient participants of the MedSeq Project. We received funding through grants U01-HG006500, U41-HG006834 and R01-HG005092 from the National Human Genome Research Institute and grant U19-HD077671 from the National Institute of Child Health and Human Development, both of the NIH. The NIH had no role in the study design; collection, analysis, and interpretation of data; the writing of this report; or the decision to submit the paper for publication.

## Disclosure Statement

J.L.V., H.L.M., C.A.M., C.E.S., D.L., J.B.K., H.L.R., and R.C.G. are employed by Partners Healthcare, which is the sole owner of GeneInsight, LLC. The authors declare that they have no other conflicts of interest.

## References

- Yang Y, Muzny DM, Reid JG, Bainbridge MN, Willis A, Ward PA, Braxton A, Beuten J, Xia F, Niu Z, Hardison M, Person R, Bekheirnia MR, Leduc MS, Kirby A, Pham P, Scull J, Wang M, Ding Y, Plon SE, Lupski JR, Beaudet AL, Gibbs RA, Eng CM: Clinical whole-exome sequencing for the diagnosis of mendelian disorders. *N Engl J Med* 2013;369:1502–1511.
- Jacob HJ, Abrams K, Bick DP, et al: Genomics in clinical practice: Lessons from the front lines. *Sci Transl Med* 2013;5:194–195.
- Green RC, Biesecker LG: Diagnostic clinical genome and exome sequencing. *N Engl J Med* 2014;370:2418–2425.
- Egan JB, Shi CX, Tembe W, Christoforides A, Kurdoglu A, Sinari S, Middha S, Asmann Y, Schmidt J, Braggio E, Keats JJ, Fonseca R, Bergsagel PL, Craig DW, Carpten JD, Stewart AK: Whole-genome sequencing of multiple myeloma from diagnosis to plasma cell leukemia reveals genomic initiating events, evolution, and clonal tides. *Blood* 2012;120:1060–1066.
- Dawson SJ, Tsui DW, Murtaza M, Biggs H, Rueda OM, Chin SF, Dunning MJ, Gale D, Forsheo T, Mahler-Araujo B, Rajan S, Humphray S, Becq J, Halsall D, Wallis M, Bentley D, Caldas C, Rosenfeld N: Analysis of circulating tumor DNA to monitor metastatic breast cancer. *N Engl J Med* 2013;368:1199–1209.
- Alexandrov LB, Nik-Zainal S, Wedge DC, et al: Signatures of mutational processes in human cancer. *Nature* 2013;500:415–421.
- Duzkale H, Shen J, McLaughlin H, Alfares A, Kelly M, Pugh T, Funke B, Rehm H, Lebo M: A systematic approach to assessing the clinical significance of genetic variants. *Clin Genet* 2013;84:453–463.
- ACMG Board of Directors: Points to consider in the clinical application of genomic sequencing. *Genet Med* 2012;14:759–761.
- Bowen MS, Kolor K, Dotson WD, Ned RM, Khoury MJ: Public health action in genomics is now needed beyond newborn screening. *Public Health Genomics* 2012;15:327–334.
- Grosse SD, Rogowski WH, Ross LF, Cornel MC, Dondorp WJ, Khoury MJ: Population screening for genetic disorders in the 21st century: evidence, economics, and ethics. *Public Health Genomics* 2010;13:106–115.
- Berg JS, Amendola LM, Eng C, Van Allen E, Gray SW, Wagle N, Rehm HL, DeChene ET, Dulik MC, Hisama FM, Burke W, Spinner NB, Garraway L, Green RC, Plon S, Evans JP, Jarvik GP: Processes and preliminary outputs for identification of actionable genes as incidental findings in genomic sequence data in the clinical sequencing exploratory research consortium. *Genet Med* 2013;15:860–867.
- Vassy JL, Lautenbach DM, McLaughlin HM, et al: The MedSeq Project: a randomized trial of integrating whole genome sequencing into clinical medicine. *Trials* 2014;15:85.
- Richards CS, Bale S, Bellissimo DB, Das S, Grody WW, Hegde MR, Lyon E, Ward BE: ACMG recommendations for standards for interpretation and reporting of sequence variations: revisions 2007. *Genet Med* 2008;10:294–300.
- Rehm HL, Bale SJ, Bayrak-Toydemir P, Berg JS, Brown KK, Deignan JL, Friez MJ, Funke BH, Hegde MR, Lyon E: ACMG clinical laboratory standards for next-generation sequencing. *Genet Med* 2013;15:733–747.
- Nippert I, Harris HJ, Julian-Reynier C, Kristofferson U, Ten Kate LP, Anionwu E, Benjamin C, Challen K, Schmidtke J, Nippert RP, Harris R: Confidence of primary care physicians in their ability to carry out basic medical genetic tasks – a European survey in five countries – part 1. *J Community Genet* 2011;2:1–11.
- Selkirk CG, Weissman SM, Anderson A, Hurlick PJ: Physicians' preparedness for integration of genomic and pharmacogenetic testing into practice within a major healthcare system. *Genet Test Mol Biomarkers* 2013;17:219–225.

- 17 Korf BR, Berry AB, Limson M, Marian AJ, Murray MF, O'Rourke PP, Passamani ER, Relling MV, Tooker J, Tsongalis GJ, Rodriguez LL: Framework for development of physician competencies in genomic medicine: report of the competencies working group of the inter-society coordinating committee for physician education in genomics. *Genet Med* 2014;16:804–809.
- 18 Haga SB, Burke W, Ginsburg GS, Mills R, Agans R: Primary care physicians' knowledge of and experience with pharmacogenetic testing. *Clin Genet* 2012;82:388–394.
- 19 Lane W, Leshchiner I, Boehler S, Uy J, Agud M, Smeland-Wagman R, Green R, Rehm H, Kaufman R, Silberstein L, Project fTM: Comprehensive blood group prediction using whole genome sequencing data from the MedSeq Project (abstract). Proceedings of the 63rd Annual Meeting of the American Society of Human Genetics, Boston, October 22–26, 2013.
- 20 Whirl-Carrillo M, McDonagh EM, Hebert JM, Gong L, Sangkuhl K, Thorn CF, Altman RB, Klein TE: Pharmacogenomics knowledge for personalized medicine. *Clin Pharmacol Ther* 2012;92:414–417.
- 21 Korf BR, Ledbetter D, Murray MF: Report of the Banbury Summit Meeting on the evolving role of the medical geneticist, February 12–14, 2006. *Genet Med* 2008;10:502–507.
- 22 Aronson SJ, Clark EH, Babb LJ, Baxter S, Farwell LM, Funke BH, Hernandez AL, Joshi VA, Lyon E, Parthum AR, Russell FJ, Varugheese M, Venman TC, Rehm HL: The GeneInsight Suite: a platform to support laboratory and provider use of DNA-based genetic testing. *Hum Mutat* 2011;32:532–536.

## Erratum

---

In the article by Jason L. Vassy et al., entitled ‘A one-page summary report of genome sequencing for the healthy adult’ [Public Health Genomics 2015;18:123–129, DOI: 10.1159/000370102], the name of the second author is wrong. The correct name is Heather M. McLaughlin.