



ARTICLE

Operationalizing the Wilson-Jungner principles for the genomics era: Consensus recommendations from the International Consortium on Newborn Sequencing



Lilian Downie^{1,2}, Julie Yeo³, Thomas Minten⁴, Rose Heald^{5,6}, Derek Ansel⁷, Mei Baker⁸, Jorune Balciuniene⁹, Jonathan S. Berg¹⁰, François Boemer¹¹, Wendy K. Chung^{12,13}, Heidi L. Cope¹⁴, David J. Eckstein¹⁵, Nicolas Encina^{6,16}, Laurence Faivre^{17,18}, Alessandra Ferlini¹⁹, Judit García-Villoria^{20,21,22}, Michael H. Gelb²³, José Manuel González De Aledo-Castillo²⁰, Katie Golden-Grant²⁴, Richard B. Parad^{13,25}, Nidhi Shah^{26,27}, Zornitza Stark^{1,2}, Kristen L. Sund²⁸, Petros Tsipouras²⁹, Meekai To³⁰, David Bick³⁰, Robert C. Green^{6,13,31,32}, International Consortium on Newborn Sequencing, Nina B. Gold^{3,33,*}

ARTICLE INFO

Article history:

Received 9 April 2025
 Received in revised form
 3 October 2025
 Accepted 9 October 2025
 Available online 24 October 2025

Keywords:

Delphi technique
 Gene selection
 Genomic sequencing
 Newborn screening

ABSTRACT

Purpose: For decades, the selection of disorders included in newborn screening (NBS) programs has been guided by principles published by Wilson and Jungner in 1968. As research explores the expansion of conditions included in NBS through genomic sequencing, there is a critical need for updated recommendations to address the opportunities and complexities of genomic data.

Methods: The International Consortium on Newborn Sequencing includes leaders from over 16 research projects investigating genomic NBS across the United Kingdom, Europe, United States, and Oceania. Consortium members were invited to participate in a modified Delphi study, aggregating opinion on the selection of conditions for genomic NBS through 3 rounds of online questionnaires, with feedback provided to participants between rounds.

Results: In round 1, 94 participants completed the questionnaire, and 10 of 43 statements reached consensus. In round 2, 81 participants completed the questionnaire, and 14 of 27 statements reached consensus. In round 3, 68 participants completed the questionnaire, and all 10 statements reached 72% or more consensus.

Conclusion: The 10 consensus recommendations developed in this study can guide future research and public health programs performing genomic NBS. This process also identified key areas of participant discordance, highlighting important topics for future research.

Published by Elsevier Inc. on behalf of American College of Medical Genetics and Genomics.

The names of the International Consortium on Newborn Sequencing members will appear at the end of the article.

*Correspondence and requests for materials should be addressed to Nina B. Gold, Mass General Hospital for Children, Division of Medical Genetics and Metabolism, 175 Cambridge Street, Boston, MA 02114. *Email address:* ngold@mgh.harvard.edu

Affiliations are at the end of the document.

doi: <https://doi.org/10.1016/j.gim.2025.101618>

1098-3600/Published by Elsevier Inc. on behalf of American College of Medical Genetics and Genomics.

Introduction

Newborn screening (NBS), introduced in the 1960s, is a successful public health intervention that identifies infants at risk for treatable disorders.¹ For nearly 6 decades, selection of disorders included in NBS programs has been guided by 10 principles published by James Wilson and Gunnar Jungner in 1968, which emphasize the inclusion of infant-onset, treatable disorders ([Supplemental Table 1](#)).² Based on these principles, the United States Health Resources and Services Administration's Recommended Uniform Screening Panel suggests that 38 core and 26 secondary conditions be screened by each state NBS program, and other countries have similar lists.³ As of 2025, over 800 genetic disorders have treatments and are potential candidates for early detection.⁴ Many such disorders cannot be identified through biochemical testing, which has led to growing interest in using genomic sequencing to expand NBS.¹ Genetic information has already been incorporated into population NBS as a first-tier screening test in several jurisdictions,^{5,6} and several international research studies have explored the feasibility of expanding genomic newborn screening (gNBS) to include hundreds or thousands of other disorders (Boemer F, Hovhannesian K, Piazzon F, et al. Population-based, first-tier genomic newborn screening in a single maternity ward in Belgium: results of babydetect project. *Social Science Research Network*. <https://doi.org/10.2139/ssrn.4896054>),⁷⁻¹⁵ with more studies underway.¹⁶⁻¹⁹ Parents,²⁰ rare disease specialists,²¹ primary care physicians,²² genetic counselors,^{21,23} and the public²⁴ support the implementation of gNBS for wider detection of disorders similar to those already screened, which are largely metabolic and endocrine, severe, and treatable. A proportion also supports inclusion of conditions for which treatment is limited or supportive. The majority value accuracy of the test over actionability.²⁵

Over 30 newborn and pediatric research programs and companies offering gNBS have been developed. Although the Wilson-Jungner principles have been utilized as a guide for implementing these studies,²⁶ they are not applicable to complex genomic information, prompting a need for additional principles to be developed.²⁷ For instance, some genes can lead to multiple phenotypes, or a spectrum of severity, age of onset, or penetrance making it unclear if they should be included for screening.²⁸ Among gNBS programs, there has been tension between maximizing the capabilities of genomic sequencing by screening for thousands of conditions, versus a conservative approach to align with the public health NBS context. This has resulted in limited concordance on which genes and variants should be analyzed and reported.²⁹⁻³¹ As gNBS moves from a research investigation to a public health approach, updated guidelines are needed to identify appropriate disorders for population screening.

The International Consortium on Newborn Sequencing (ICoNS) is a professional organization founded in 2022. It aims to aggregate data from gNBS studies to best

implement their findings. The ICoNS gene list subcommittee, which developed this study, was created with the aim of defining principles of gene and variant selection for public health gNBS. The aim of this study was to define principles to guide gene selection for genomic newborn screening in a population health screening context. Other pertinent issues related to gNBS such as universality, equity, accessibility, cost, follow-up, consent, and data storage were outside the scope of this work and are under active investigation by other subcommittees within ICoNS.

Materials and methods

Study design

The Delphi method provides a transparent process by which a group of experts can reach consensus.³² It involves iterative rounds of anonymized questionnaires with aggregated results presented between rounds to enable individuals to reflect on collective opinion and refine their own position. Methodology was informed by the Recommendations for the Conducting and REporting of DELphi Studies.³³ Modifications were incorporated to accommodate the participation of a large group of experts and stakeholders located in a broad geographical area. The study was conducted online and included variation in the participants for each round, in contrast to the traditional selection of a fixed group of experts.

Participants

ICoNS membership is available to any individual that identifies as a stakeholder in gNBS. Members are required to disclose their position, expertise and interest in gNBS and financial or conflicting interests at the time of application. Membership falls into the categories of

- Clinicians involved in genomics or newborn screening. These include clinical geneticists, genetic counselors and physicians. This group includes leaders from each of the large projects investigating gNBS across the United Kingdom, Europe, the United States, and Oceania ([Figure 1](#)).
- Scientists working in both genomics and in traditional newborn screening methods.
- Researchers, including academics from multiple disciplines (medicine, health economics, implementation science, etc) with an interest or position in a genomic newborn sequencing study.
- Industry, which includes those employed by companies with an interest in genomics, newborn screening or precision medical therapies. This group includes those with financial and commercial interest in gNBS. The consortium actively chose to be inclusive of this group because some are already offering direct to consumer




Figure 1 Map of global gNBS research programs represented in ICoNS. Countries of research programs are indicated in parentheses. Intended enrollment sizes are indicated where available. Shading on map refers to the locations of individual ICoNS members.

gNBS. This was seen to be mutually beneficial because those with a commercial interest may provide expertise on scaling and automation required to provide population level screening. The consortium membership in return can provide ethical and evidence-based guidance on implementation.

At the time the first Delphi questionnaire was circulated (June 25, 2024), ICoNS had 234 registered members across 28 countries. By the second Delphi questionnaire, the consortium had expanded to 256 members across 28 countries and 27 US states (August 8, 2024) and grew to 270 members across 32 countries and 28 US states by the final Delphi questionnaire (September 16, 2024). The breakdown of membership is 85% clinicians, scientists, and researchers and 15% from industry.

The gene list subcommittee was established in February 2024 and comprises 25 members from 8 countries. The group met monthly via Zoom (March 19, 2024; April 23, 2024; May 28, 2024; July 23, 2024; and August 27, 2024) to plan and design the study. The composition of this group was by self-nomination and by invitation with an aim to ensure representation across the different stakeholder groups.

Questionnaire development

A literature review of criteria for gNBS public health and research programs was completed (Supplemental Table 2). These criteria, along with topics generated by the gene list subcommittee, were utilized to develop a set of 43 statements

pertaining to the following categories: age of symptom onset and age of actionability (7 statements), prevalence and gene-disease validity (3 statements), penetrance (5 statements), clinical features of disease (5 statements), variant reporting (8 statements), variant calling and technical interpretation (5 statements), nongenetic confirmatory testing (5 statements), treatment (4 statements), and parental engagement (1 statement). Three to 4 gene list subcommittee members were tasked with refining statements within each category. The final set of 43 statements were developed into a questionnaire using an online survey tool (REDCap) (Supplemental Note 1).^{34,35}

For most statements, participants were offered only “agree” or “disagree” response options to encourage clear answers. For some statements, checkboxes that allowed for multiple selections, a slider with continuous variables, or radio buttons were offered. All questions required a response to continue to the next section. An optional area for free-text responses was included at the end of each category, allowing participants to provide comments, suggestions, or clarifications on their responses.

Per the modified Delphi model, results of the round 1 questionnaire were collated and presented back to ICoNS members via an online presentation and email. Statements that reached the consensus threshold were integrated into the statements presented in the round 3 questionnaire (Figure 2). All other statements and free-text responses were utilized to refine, remove, or add questions for the round 2 questionnaire (Supplemental Note 2). Statements from the round 2 questionnaire that reached consensus were




Figure 2 Delphi process for updating gene and disorder selection criteria for population-wide genomic newborn screening. Statements related to gNBS gene and disorder selection criteria were distributed through 3 rounds of questionnaires. Consensus statements from the first 2 rounds were compiled to produce 10 consensus guidelines.

combined with those from round 1 and presented as a round 3 questionnaire (Supplemental Note 3).

Distribution of questionnaire

The questionnaire was sent by email to all ICoNS members. Participants were asked to consider the statements in the context of public health gNBS, not research programs. Participants were not asked to disclose any identifying data, making all responses anonymous. For this reason, approval from the Institutional Review Board was not required. The questionnaire was piloted before distribution with 4 individuals who had expertise in medical genetics. These responses were not included in the analysis.

Definition of consensus

There is variation in the literature about what proportion of responses constitutes consensus from the Delphi process.³⁶ For this process, consensus was defined as $\geq 85\%$ participant agreement for round 1, $\geq 75\%$ for round 2, and $\geq 70\%$ for round 3. Approaching consensus was defined as $\geq 60\%$ and divided opinion as 41% to 59% agreement. The threshold for consensus was set very high in the initial round to efficiently identify areas of high agreement or disagreement and focus on more contentious topics to get the highest value out of the process.

Results

Round 1 questionnaire

In round 1, 94 completed questionnaires were received (40.2% response rate). Ten statements reached consensus, 25 statements were approaching consensus (60%-85% consensus), and the remaining 8 statements demonstrated divided opinion below 60% agreement (Figure 3). The statements with the highest consensus included: “Age of symptom onset is an important factor in gene list selection” (96% of participants agreed [$n = 91$]) and “Genes where most disease causing variants are inherited and highly penetrant (for example, deleterious variants in *FBNI* [HGNC:3603], which typically lead to Marfan syndrome [MIM:154700]) should be included” (95% of participants agreed [$n = 89$]) (Supplemental Table 3).

Some statements that were approaching consensus were accompanied by free-text responses that allowed the statement to be refined, for example: “Some disorders for which there is not a confirmatory, non-genetic test should be included” (81% of participants agreed [$n = 70$]) with free-text statements, such as: “If there is no confirmatory test but you are quite certain about gene-disease validity and a life-saving treatment needs to be administered quickly, confirmatory testing may not be necessary.” Eight statements that were either approaching consensus or divided were removed based on feedback from free-text responses or because of overlap with other recirculated statements. Two new statements were added after reviewing the free-text responses: “Variants associated with disorders that manifest symptoms and require treatment or surveillance only in adulthood (for example, hereditary breast and ovarian cancer): A. Should be reported because a positive result may improve the health of a parent. B. Should be reported because they are relevant to the child’s future health. C. Should not be reported.” and “Variants in genes such as *BRCA1* (HGNC:1100), which can cause an autosomal recessive childhood disorder, should only be reported when homozygous or biallelic, even if the monoallelic disease has implications for health.”

The most discordant results were in the category of penetrance. For example, “All likely pathogenic and pathogenic variants should be included in newborn screening programs even if the penetrance is not known” (53% of participants agreed [$n = 50$]) and “Low penetrance genotypes may be included for diseases in which the treatment is inexpensive, readily available, and of minimal risk to the well-being of the patient (such as *MYH7* (HGNC:7577)-related cardiomyopathy where echocardiogram surveillance may be recommended)” (57% of participants agreed [$n = 54$]). Seven out of 8 of the divided statements were recirculated in the round 2 questionnaire, and 1 statement, “All likely pathogenic and pathogenic variants should be included in newborn screening programs even if the penetrance is not known,” was removed because of overlap with other recirculated statements.




Figure 3 Summary of consensus levels across Delphi rounds.

A total of 43 statements were distributed in the round 1 questionnaire across 10 different categories related to gNBS. A total of 27 statements were circulated in the round 2 questionnaire.

A total of 59 participants submitted 238 free-text responses, which included rationales for the answers selected by participants, feedback on the quality of the question and its wording, and suggestions for future questions. Themes that emerged from the free-text responses in the round 1 questionnaire included the importance of considering the questions for a public health screening context, limited data about penetrance, and the nuances for each gene-disease pair that increase the challenge of creating universal guidelines.

The statement that generated the highest number of free-text comments was: “Parents should be given an opportunity to choose which genes are screened in their child.” In total, 54 participants (64%) disagreed with this statement. Free-text comments in response to this statement included: “an option for parents to choose would create havoc” and “parents cannot be expected to make decisions about what genes to screen.” Multiple free-text responses suggested utilizing categories of disease to provide parents’ a more informed choice. These responses included: “if there is choice it should not be on the gene level but perhaps they can opt in for ‘less actionable’ conditions as a group” and “parents can make choices in classes of genes in relation to disease severity and treatment options.”

Round 2 questionnaire

In round 2, 81 participants completed the questionnaire (31.6% response rate). Of 27 statements, 14 reached

consensus, 7 were approaching consensus, and the remaining 6 demonstrated divided opinion (Supplemental Table 4). There were 159 free-text responses from 42 participants.

Several statements that were approaching consensus in round 1 achieved consensus in round 2. These included: “If there are well-established genotype-phenotype correlations between variants in a gene and the onset of disease (for example, in lysosomal acid lipase deficiency [MIM:620151]), only variants associated with the childhood-onset form of the condition should be included” (75% of participants agreed [n = 60]), “Different sets of genes or variants should be queried at various ages throughout a person’s lifespan” (80% of participants agreed [n = 64]), “If meeting all other inclusion criteria, highly prevalent monogenic disorders (such as familial hypercholesterolemia [MIM:143890], G6PD deficiency [MIM:300908]) should be included in genomic newborn screening, even though it would result in increased follow-up burden for health care providers” (87% of participants agreed [n = 69]), and “Moderate penetrance genotypes may be reported if there is nongenetic confirmatory testing for the disease available. (For example, variants in *HFE* [HGNC:4886], which may cause hemochromatosis [MIM:235200], can be adjudicated with laboratory tests such as a serum ferritin and transferrin saturation)” (89% of participants agreed [n = 66]).

Among the 8 statements with divided opinion from round 1 that were recirculated in round 2, one reached consensus in round 2: “Variants associated with mild differences in body structure and/or function should be included (for example, variants in *UGT1A1* [HGNC:12530] related to Gilbert syndrome [MIM:143500])” (40% of participants agreed in round 1 [n = 38], this fell to 14% agreement in round 2). To summarize, 86% [n = 64] of participants thought screening for mild conditions should not be included. Six statements with divided opinion in round 1 had responses that were approaching consensus in round 2. Only 1 statement that demonstrated divided opinion in round 1 continued to demonstrate divided opinion in round 2: “For genes associated with recessive disorders, if a single pathogenic or likely pathogenic variant is identified, the gene should also be queried for variants of uncertain significance and these should be reported” (50% of participants agreed in round 1 [n = 45], 42% of participants agreed in round 2 [n = 31]) (Table 1).

Free-text responses in round 2 highlighted additional themes, some of which were not observed in round 1. For example, several participants noted that health care resources should not be a significant factor in determining which conditions to screen because each country’s health care system will adapt over time and reallocate care as needed, “[W]e should not let the burden of health care providers stop us from doing the right thing for NBS. Such burden can be short term (i.e. in the long term, care burden for G6PD patients would decrease), and there are other ways to address resource issues if well anticipated.”

Table 1 Statements across Delphi rounds with divided opinion responses (<60% consensus)

Statement	Round	Suggested Areas of Future Research
Age of symptom onset and age of actionability If the earliest reported case of a predominantly adult disorder is in childhood, then the associated disease gene should be included. (For example, PMID: 36313796 describes four cases of children who were diagnosed with Lynch syndrome-associated colorectal cancers age 14-17 years of age.)	Round 1	Review of literature for disorders with a large range of ages of onset
Prevalence and gene-disease validity If meeting all other criteria for inclusion, genes with gene-disease validity of moderate or higher based on ClinGen gene-disease validity scoring criteria should be included on genomic newborn screening.	Round 1	Additional curation of gene-disease relationships
Penetrance All likely pathogenic and pathogenic variants should be included in newborn screening programs even if the penetrance is not known.	Round 1	Studies of biobanks and large data sets and longitudinal follow-up of infants with positive gNBS results
Low penetrance genotypes may be included for diseases in which the treatment is inexpensive, readily available, and of minimal risk to the well-being of the patient (such as MYH7 related cardiomyopathy where echocardiogram surveillance may be recommended).	Round 1	
Only variants with well-established penetrance estimates should be reported in newborn screening programs.	Round 2	
Clinical features of disease Genes associated with phenotypes that cause mild differences in body structure and/or function should be included, for example nonclassical congenital adrenal hyperplasia.	Round 1	Development of a health instrument to measure the severity of genetic disease
Variant curation and reporting For genes associated with recessive disorders, if a single pathogenic or likely pathogenic variant is identified, then the gene should also be queried for variants of uncertain significance, and these should be reported.	Round 1	Studies of biobanks and large data sets to assess sensitivity and specificity of including variants of uncertain significance
For genes associated with recessive disorders, if a single pathogenic or likely pathogenic variant is identified, then the gene should also be queried for variants of uncertain significance, and these should be reported.	Round 2	
Heterozygous variants associated with X-linked disorders should be reported when found in female infants when they are known to be associated with any phenotype, even if that phenotype is more mild or later onset (ie, G6PD deficiency).	Round 1	Deep phenotyping of females with variants associated with X-linked disorders
Treatment Monogenic disorders should only be selected for inclusion in genomic newborn screening if the appropriate treatment is available in the country in which screening is being performed.	Round 1	Global assessment of availability of targeted therapies for genetic disorders
Nongenetic confirmatory testing A monogenic condition should be eligible for genomic newborn screening as long as a nongenetic confirmatory test is expected to become abnormal before treatment is initiated.	Round 2	Studies of penetrance and development of additional biomarkers of disease
Parental engagement Parents should be given an opportunity to choose if their child is screened for certain types of disorders (for example, treatable or nontreatable).	Round 2	Qualitative research with parents and people with genetic conditions

G6PD, glucose-6-phosphate dehydrogenase; *gNBS*, genomic newborn screening.

A total of 24 statements reached consensus across rounds 1 and 2. Two additional statements were within 1% of consensus in round 2. Statements which were beyond the scope of gene list curation ($n = 5$), for example, those regarding the reanalysis of genomic data across the lifespan, were not included in the round 3 questionnaire. Statements that contained repetitive information or granular detail were refined, for example, those relating to carrier status for recessive conditions and adult-onset symptoms in individuals with heterozygous variants in autosomal recessive disease genes were combined into 1 statement.

Round 3 questionnaire

The remaining 19 statements that had achieved consensus in the prior 2 rounds were combined and edited by the subcommittee leaders to produce 10 final statements, which were circulated as the round 3 questionnaire (Table 2). Examples of combined statements include those relating to nonconfirmatory genetic testing and penetrance (Supplemental Table 4), in which 4 statements reached consensus and were readily combined into a single statement “Variants with incomplete penetrance should only be reported if there is a nongenetic confirmatory test expected to be positive before initiation of treatment, or if the disease surveillance or management presents minimal risk.” Statements that had not achieved consensus by round 2 were not included in round 3. In round 3, 68 participants completed the questionnaire (25.2% response rate), and all statements reached 72% or greater levels of consensus. Statements were compared with the Wilson and Juenger principles (Table 3).

Discussion

This study presents 10 statements based on expert consensus that provide detail and guidance on the selection of conditions and reporting of variants for gNBS programs, which may be

used to guide future global public health programs. Although the Wilson-Jungner principles² have historically provided a framework by which to choose disorders for NBS, the growing number of targeted therapies for genetic disorders and the complexities of genomic information provide an opportunity to consider a new, more nuanced set of recommendations.^{4,26} Studies have demonstrated that gNBS has the potential to improve detection rates of genetic disease in apparently healthy infants,³⁷ without causing undue psychosocial harm or damage to infant-parent bonding.³⁸ However, selection of appropriate genes, conditions, and variants will be critical to the successful implementation of gNBS as a public health approach. Successful implementation will depend not solely on detection rate but on overall health benefit for individuals and families. Participants’ consensus centered around several key themes. Participants indicated that gNBS, similar to NBS, performed through biochemical methods, should continue to focus on severe diseases that are urgent and treatable with intervention in early childhood. This reflects findings from another recent Delphi focused on treatability in the context of gNBS, which concluded that the expected benefit/burden ratio of early treatment should be positive and result in a significant health outcome.³⁹ Importantly, emphasis was placed not on the time when symptoms were expected to emerge but when surveillance or treatment for the disorder might begin; for example, Alport syndrome (MIM: 301050, 203780, 620536, and 104200) typically leads to adult-onset kidney failure and hearing loss, but antihypertensive medications are recommended for use in early childhood to prevent disease progression.⁴⁰ Also aligned with Wilson and Jungner, who stated that uncommon disorders could be screened so long as “very serious consequences [were expected] if [the disease was] not discovered and treated very early in life,” these guidelines emphasize that the prevalence of the disease is not relevant, particularly when many conditions can be screened simultaneously.² However, in contrast to the Wilson-Jungner principles, which states that there “should be a suitable test or examination” to confirm the presence of disease, participants indicated that a nonmolecular diagnostic test does not

Table 2 International Consortium on Newborn Sequencing (ICoNS) consensus guidelines for gene selection in genomic newborn screening programs

Consensus Statements

1. Variants should be associated with a disease for which treatment or surveillance should be initiated before age 5.
2. Disorders should have a treatment that greatly improves the severity of disease.
3. Variants associated with mild differences in body structure and/or function should not be reported.
4. Variants should be included, even if the associated disorder is unlikely to escape detection by a clinical team.
5. Sequencing of the infant’s sample alone is a sufficient sequencing approach for genomic newborn screening.
6. Variants with incomplete penetrance should only be reported if there is a non-genetic confirmatory test expected to be positive before initiation of treatment, or if the disease surveillance or management presents minimal risk.
7. Genes in which the well described variants cannot easily be detected on sequencing should be included in genomic newborn screening, because other variants may be ascertained.
8. Disease prevalence should not be a criteria for variant reporting in a population.
9. Carrier status of recessive disorders should not be reported.
10. If only variants of uncertain significance are found in a gene, no variants should be reported.

Table 3 International Consortium on Newborn Sequencing (ICoNS) consensus guidelines for gene selection in genomic newborn screening programs as compared with the original Wilson-Jungner principles

ICoNS Consensus Statements	Corresponding Wilson-Jungner Principle(s)	Implications for gNBS
<ol style="list-style-type: none"> 1. Variants associated with mild differences in body structure and/or function should not be reported. 2. Disease prevalence should not be a criteria for variant reporting in a population. 3. Carrier status of recessive disorders should not be reported. 	<ol style="list-style-type: none"> 1. The condition should be an important health problem. 	<p>Wilson and Jungner wrote that rare disorders, such as PKU, are suitable for screening due to the severe consequences of delayed treatment.¹ ICoNS members agreed that the prevalence of a genetic condition should not determine its inclusion in gNBS.</p> <p>ICoNS members also agreed that genetic variants linked to mild conditions, such as Gilbert syndrome, and reproductive risks without personal medical implications, like carrier status for AR disorders, should be excluded. Importantly, ICoNS reached consensus that AD variants associated with adult-onset disorders in genes that are also linked to AR childhood-onset disorders (eg, BRCA1, which causes both AD HBOC and AR Fanconi anemia) should not be reported on gNBS.</p>
<ol style="list-style-type: none"> 4. Disorders should have a treatment that greatly improves the severity of disease. 	<ol style="list-style-type: none"> 2. There should be an accepted treatment for patients with recognized disease. 3. Facilities for diagnosis and treatment should be available. 	<p>ICoNS members agreed that there should be a treatment available, and that a nongenomic confirmatory test is not required for a gene to be suitable for screening.</p> <p>ICoNS members did not reach consensus on whether a condition should be included in gNBS if the appropriate treatment is unavailable in the country in which screening is being performed.</p>
<ol style="list-style-type: none"> 5. Sequencing of the infant’s sample alone (eg, no parental DNA samples needed) is a sufficient sequencing approach for genomic newborn screening. 6. Genes in which the well described variants cannot easily be detected on sequencing should be included in genomic newborn screening, because other variants may be ascertained. 	<ol style="list-style-type: none"> 4. There should be a recognizable latent or early symptomatic stage. 5. There should be a suitable test or examination. 6. The test should be acceptable to the population. 7. The natural history of the condition, including development from latent to declared disease, should be adequately understood. 	<p>A latent or early symptomatic stage was not considered necessary for many genetic disorders to be included in gNBS. ICoNS members agreed that sequencing the infant’s sample alone, rather than duo or trio sequencing with parents, is sufficient and practical. For some conditions, such as hemophilia A, technologies like ES may fail to detect common genetic changes underlying the disease (eg, inversions). Despite this limitation, ICoNS members agreed that these genes should still be included in gNBS to identify individuals with other P or LP variants.</p> <p>The natural history of most genetic disorders is poorly understood, particularly because they are individually rare and early research often focuses on the most severely affected individuals.</p>

(continued)

Table 3 Continued

ICoNS Consensus Statements	Corresponding Wilson-Jungner Principle(s)	Implications for gNBS
<p>7. Variants should be associated with a disease for which treatment or surveillance should be initiated before age 5.</p> <p>8. Variants with incomplete penetrance should only be reported if there is a non-genetic confirmatory test expected to be positive before initiation of treatment, or if the disease surveillance or management presents minimal risk.</p> <p>9. If only variants of uncertain significance are found in a gene, no variants should be reported.</p> <p>10. Variants should be included, even if the associated disorder is unlikely to escape detection by a clinical team.</p>	<p>8. There should be an agreed policy on whom to treat as patients.</p> <p>9. The cost of case finding should be economically balanced in relation to possible expenditure on medical care as a whole.</p> <p>10. Case finding should be a continuing process and not a “once and for all” process.</p>	<p>ICoNS members agreed that the existence of an orthogonal, nonmolecular confirmatory test was unnecessary for inclusion in gNBS, except in cases where the variant or disorder is known to have incomplete penetrance. They emphasized that as long as at least 1 variant in a gene is a P or LP (eg, not 2 VUS in a gene with AR inheritance) and is associated with a condition requiring treatment or surveillance before age 5, it should be included. Additionally, ICoNS members agreed that variants linked to disorders such as achondroplasia, which present with physical findings at birth, should still be included.</p> <p>A separate ICoNS subcommittee is dedicated to assessing the economic considerations related to gNBS.</p> <p>ICoNS members did reach consensus on the statement, “Different sets of genes or variants should be queried at various ages throughout a person’s lifespan,” but this area was considered beyond the scope of gene selection for gNBS.</p>

AD, autosomal dominant; *AR*, autosomal recessive; *ES*, exome sequencing; *gNBS*, genomic newborn screening; *HBOC*, hereditary breast and ovarian cancer; *ICoNS*, International Consortium of Newborn Sequencing; *LP*, likely pathogenic; *P*, pathogenic; *PKU*, phenylketonuria.

necessarily need to be available after a positive gNBS result and before treatment initiation if the associated treatment is low-risk or inexpensive. This suggestion that genetic information is in some cases sufficient to guide care would allow expansion of gNBS to a wider range of disorders, including hereditary cancer predisposition syndromes affecting young children or neurodevelopmental disorders that may not be symptomatic in infancy.

This premise needs to be balanced with the potential harm of medicalizing a healthy infant and the burden that anticipatory care can place on families. This is recognized by the consensus that mild conditions, even if they can be readily detected, should still be excluded from newborn screening programs. It also provides the basis for the suggestions for future research in which penetrance and expression data are going to be key to responsible reporting of variants in future programs.

Participants demonstrated divided opinions on a range of topics, which are important areas for future research. Several of the categories are specific to genomic information, for example, the reporting of variants of uncertain significance when found in the context of a biallelic pathogenic (P) or likely pathogenic (LP) variant. Ongoing gNBS studies approach this challenge differently; for example, in the GUARDIAN study,¹¹ when infants with a single P or LP variant in a gene associated with autosomal recessive inheritance are identified, their samples are also evaluated for some variants of uncertain significance in the same gene, and second-tier nonmolecular orthogonal testing is performed when available. The downstream sensitivity and specificity of this approach, compared with the reporting of P or LP variants alone, is an important area of investigation. One long-term challenge highlighted by this study is the lack of accurate data regarding disease penetrance and expressivity. Because individuals with most genetic conditions have historically been identified after the onset of symptoms, the individual risk to develop phenotypic manifestations often remains unknown based solely on the variants identified. Currently, studies of individuals at risk for genetic disease from biobanks and large data sets may provide insights into penetrance; however, most of these biobanks are exclusively composed of adults.^{28,41-43} Furthermore, given the large number of different combinations of alleles for biallelic conditions, there may not be empirical data to guide a particular combination observed for the first time in gNBS. The medical outcomes of infants from gNBS pilot studies will provide additional longitudinal information in this area.³⁷

All elements of gNBS introduction, such as health economics, the impact on care pathways and evolution of precision planning, and the impact on individual families require significantly larger studies to generate evidence. These studies are underway, and funding bodies have recognized the need for these data to implement responsible and acceptable gNBS programs. These guidelines are timely to provide guidance on how to select conditions and variants to report in this context.

Limitations to this study include a modification of the classic Delphi process. This was chosen to maximize participation from a large group of experts and stakeholders. Typically, in a Delphi model, the same participants are selected for their expertise and invited in each round. The group chose to modify this and distribute questionnaires to all ICoNS participants for each round. This meant that the group was dynamic because the membership grew over the course of the study, and perhaps in line with this, the participation rate was lower than expected for a standard Delphi methodology. The diversity of membership, including those from industry, is viewed as a strength by the study team but could be viewed as a conflict of interest. Being inclusive of the entire membership, provided representation from a wide range of skills and perspectives but not necessarily all relevant parties. Given that gNBS is a new area of research and the implications are relevant to the whole population, preference was given to include as many voices as possible rather than a smaller group of potential experts.

gNBS holds great promise to improve the health of newborns and children but is a new frontier that requires an updated version of the guidelines that have been used in population screening in the past. The 10 recommendations developed in this study can guide further efforts in this area, including more granular guidance and approaches to building consensus on national and international scale.

Data Availability

The published article includes all data sets generated or analyzed during this study.

Funding

No funding was obtained specifically for this study. Authors' contributions were supported by the following grants: R01 HG012271 (J.S.B.); R01 DK067859 (M.G.); HD077671 and TR003201 (R.C.G); and K08HG012811 and TR003201 (N.B.G). W.K.C. receives funding support from Sanofi.

Author Contributions

Conceptualization: D.A., M.B., J.B., J.S.B., F.B., W.K.C., H.L.C., L.D., D.J.E., N.E., L.F., A.F., J.G., M.H.G., N.B.G., J.M.G.D.A.-C., K.G.-G., R.B.P., N.S., Z.S., K.L.S., P.T., M.T., J.Y.; Data Curation: L.D., N.B.G., R.H., J.Y.; Formal Analysis: L.D., N.B.G., J.Y.; Funding Acquisition: N.E.; Investigation: D.A., M.B., J.B., J.S.B., F.B., W.K.C., H.L.C., L.D., D.J.E., N.E., L.F., A.F., J.G., M.H.G., N.B.G., J.M.G.D.A.-C., K.G.-G., R.B.P., N.S., Z.S., K.L.S., P.T., M.T., J.Y.; Methodology: D.A., M.B., J.B., J.S.B., F.B., W.K.C., H.L.C., L.D., D.J.E., N.E., L.F., A.F., J.G., M.H.G.,

N.B.G., J.M.G.D.A.-C., K.G.-G., R.B.P., N.S., Z.S., K.L.S., P.T., M.T., J.Y.; Supervision: D.B., N.B.G., R.C.G.; Visualization: N.B.G., T.M., J.Y.; Writing-original draft: L.D., N.B.G., J.Y.; Writing-review and editing: all authors.

Ethics Declaration

This study involved responses from 94 participants who completed the questionnaire, and 10 of 43 statements reached consensus. Because no individualized or patient-specific data were used, Institutional Review Board (IRB) or Research Ethics Committee (REC) approval were not required.

Conflict of Interest

Derek Ansel is a board member for Gene People. Wendy K. Chung is on the board of directors for Prime Medicine. Robert C. Green receives compensation for advising the following companies: Allelica, Atria, Fabric, Genomic Life, and Juniper Genomics; and is a cofounder of Genome Medical and Nurture Genomics. Kristen L. Sund is a consultant at Nurture Genomics. Petros Tsiouras is on the board of directors for PlumCare RWE, Inc. All other authors declare no conflicts of interest.

Declaration of AI and AI-Assisted Technologies in the Writing Process

During the preparation of this work the authors used ChatGPT to check grammar and improve sentence syntax and clarity. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

Members of the International Consortium on Newborn Sequencing

See [Supplemental Material A](#) for a list of International Consortium on Newborn Sequencing (ICoNS) authors.

Additional Information

The online version of this article (<https://doi.org/10.1016/j.gim.2025.101618>) contains supplemental material, which is available to authorized users.

Affiliations

¹Victorian Clinical Genetics Services, Murdoch Children's Research Institute, Melbourne, Australia; ²Department of

Pediatrics, University of Melbourne, Melbourne, Australia; ³Division of Medical Genetics and Metabolism, Massachusetts General Hospital, Boston, MA; ⁴KU Leuven, Leuven, Belgium; ⁵Division of Genetics, Department of Medicine, Mass General Brigham, Boston, MA; ⁶Ariadne Labs, Boston, MA; ⁷Worldwide Clinical Trials, Research Triangle Park, NC; ⁸Department of Pediatrics, University of Wisconsin School of Medicine and Public Health, Madison, WI; ⁹Revvity Omics, Waltham, MA; ¹⁰The University of North Carolina at Chapel Hill, Chapel Hill, NC; ¹¹CHU Liège, University of Liège, Liège, Belgium; ¹²Department of Pediatrics, Boston Children's Hospital, Boston, MA; ¹³Harvard Medical School, Boston, MA; ¹⁴RTI International Genomics and Translational Research Center, Research Triangle Park, NC; ¹⁵Office of Clinical Research Education and Collaboration Outreach, Office of the Director, NIH, Bethesda, MD; ¹⁶Harvard T.H. Chan School of Public Health, Boston, MA; ¹⁷Genetics Center, FHU TRANSLAD, CHU Dijon Bourgogne, Dijon, France; ¹⁸UMR1231 INSERM, GAD team, Bourgogne University, Dijon, France; ¹⁹Medical Genetics Unit, Department of Medical Genetics, University of Ferrara, Ferrara, Italy; ²⁰Division of Inborn Errors of Metabolism-IBC, Department of Biochemistry and Molecular Genetics, Hospital Clinic of Barcelona, Barcelona, Spain; ²¹Center for Biomedical Research Network on Rare Diseases (CIBERER), ISCIII, Madrid, Spain; ²²Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain; ²³Departments of Chemistry and Biochemistry, University of Washington, Seattle, WA; ²⁴Rady Children's Institute for Genomic Medicine, San Diego, CA; ²⁵Department of Pediatrics, Brigham and Women's Hospital, Boston, MA; ²⁶Dartmouth Health Children's, Lebanon, NH; ²⁷Geisel School of Medicine, Hanover, NH; ²⁸Nurture Genomics, Cambridge, MA; ²⁹FirstSteps, National gNBS Initiative, Greece; ³⁰Genomics England Limited, London, England, United Kingdom; ³¹Mass General Brigham, Boston, MA; ³²Broad Institute, Boston, MA; ³³Department of Pediatrics, Harvard Medical School, Boston, MA

References

1. Stark Z, Scott RH. Genomic newborn screening for rare diseases. *Nat Rev Genet.* 2023;24(11):755-766. <http://doi.org/10.1038/s41576-023-00621-w>
2. Wilson JMG, Jungner G. *Principles and Practice of Screening for Diseases.* Geneva, Switzerland: World Health Organization; 1968.
3. Recommended Uniform Screening Panel. Accessed January 3, 2025. <https://www.hrsa.gov/advisory-committees/heritable-disorders/rusp>
4. Bick D, Bick SL, Dimmock DP, Fowler TA, Caulfield MJ, Scott RH. An online compendium of treatable genetic disorders. *Am J Med Genet C Semin Med Genet.* 2021;187(1):48-54. <http://doi.org/10.1002/ajmg.c.31874>
5. Kemper AR, Ream MA, Lam KK. Review of newborn screening implementation for spinal muscular atrophy final report. Health Resources and Services Administration (HRSA). Accessed December 9, 2024. <https://www.hrsa.gov/sites/default/files/hrsa/advisory-committees/>

- heritable-disorders/reports-recommendations/sma-nbs-implementation-report.pdf
6. Bell SC, Mall MA, Gutierrez H, et al. The future of cystic fibrosis care: a global perspective. *Lancet Respir Med.* 2020;8(1):65-124. [http://doi.org/10.1016/S2213-2600\(19\)30337-6](http://doi.org/10.1016/S2213-2600(19)30337-6)
 7. Balciuniene J, Liu R, Bean L, et al. At-risk genomic findings for pediatric-onset disorders from genome sequencing vs medically actionable gene panel in proactive screening of newborns and children. *JAMA Netw Open.* 2023;6(7):e2326445. <http://doi.org/10.1001/jamanetworkopen.2023.26445>
 8. Wang X, Sun Y, Guan XW, et al. Newborn genetic screening is highly effective for high-risk infants: a single-centre study in China. *J Glob Health.* 2023;13:04128. <http://doi.org/10.7189/jogh.13.04128>
 9. Huang X, Wu D, Zhu L, et al. Application of a next-generation sequencing (NGS) panel in newborn screening efficiently identifies inborn disorders of neonates. *Orphanet J Rare Dis.* 2022;17(1):66. <http://doi.org/10.1186/s13023-022-02231-x>
 10. Chen T, Fan C, Huang Y, et al. Genomic sequencing as a first-tier screening test and outcomes of newborn screening. *JAMA Netw Open.* 2023;6(9):e2331162. <http://doi.org/10.1001/jamanetworkopen.2023.31162>
 11. Ziegler A, Koval-Burt C, Kay DM, et al. Expanded newborn screening using genome sequencing for early actionable conditions. *JAMA.* 2025;333(3):232-240. <http://doi.org/10.1001/jama.2024.19662>
 12. Hao C, Guo R, Hu X, et al. Newborn screening with targeted sequencing: a multicenter investigation and a pilot clinical study in China. *J Genet Genomics.* 2022;49(1):13-19. <http://doi.org/10.1016/j.jgg.2021.08.008>
 13. Cao Z, He X, Wang D, et al. Targeted exome sequencing strategy (NeoEXOME) for Chinese newborns using a pilot study with 3423 neonates. *Mol Genet Genomic Med.* 2024;12(1):e2357. <http://doi.org/10.1002/mgg3.2357>
 14. Ceyhan-Birsoy O, Murry JB, Machini K, et al. Interpretation of genomic sequencing results in healthy and ill newborns: results from the BabySeq Project. *Am J Hum Genet.* 2019;104(1):76-93. <http://doi.org/10.1016/j.ajhg.2018.11.016>
 15. Shubina J, Tolmacheva E, Maslennikov D, et al. WES-based screening of 7,000 newborns: a pilot study in Russia. *HGG Adv.* 2024;5(4):100334. <http://doi.org/10.1016/j.xhgg.2024.100334>
 16. Lunke S, Bouffler SE, Downie L, et al. Prospective cohort study of genomic newborn screening: BabyScreen+ pilot study protocol. *BMJ Open.* 2024;14(4):e081426. <http://doi.org/10.1136/bmjopen-2023-081426>
 17. Peay HL, Gwaltney AY, Moultrie R, et al. Education and consent for population-based DNA screening: a mixed-methods evaluation of the Early Check newborn screening pilot study. *Front Genet.* 2022;13:891592. <http://doi.org/10.3389/fgene.2022.891592>
 18. Newborn Genomes Programme. Genomics England. Accessed January 8, 2025. <https://www.genomicsengland.co.uk/initiatives/newborns>
 19. What is Screen4Care? Accessed January 8, 2025. <https://screen4care.eu/the-project/screen4care>
 20. Gold NB, Omorodion JO, Del Rosario MC, et al. Preferences of parents from diverse backgrounds on genomic screening of apparently healthy newborns. *J Genet Couns.* 2025;34(2):e1994. <http://doi.org/10.1002/jgc4.1994>
 21. Gold NB, Adelson SM, Shah N, et al. Perspectives of rare disease experts on newborn genome sequencing. *JAMA Netw Open.* 2023;6(5):e2312231. <http://doi.org/10.1001/jamanetworkopen.2023.12231>
 22. Acharya K, Ackerman PD, Ross LF. Pediatricians' attitudes toward expanding newborn screening. *Pediatrics.* 2005;116(4):e476-e484. <http://doi.org/10.1542/peds.2005-0453>
 23. del Rosario MC, Swenson KB, Coury S, Schwab J, Green RC, Gold NB. Genetic counselors' perspectives on genomic screening of apparently healthy newborns in the United States. *Genet Med Open.* 2024;2:101885. <http://doi.org/10.1016/j.gimo.2024.101885>
 24. Lynch F, Best S, Gaff C, et al. Australian public perspectives on genomic newborn screening: risks, benefits, and preferences for implementation. *Int J Neonatal Screen.* 2024;10(1):6. <http://doi.org/10.3390/ijns10010006>
 25. Downie L, Halliday J, Lewis S, Amor DJ. Principles of genomic newborn screening programs: a systematic review. *JAMA Netw Open.* 2021;4(7):e2114336. <http://doi.org/10.1001/jamanetworkopen.2021.14336>
 26. Friedman JM. Implementing genomic newborn screening as an effective public health intervention: sidestepping the hype and criticism. *NPJ Genom Med.* 2024;9(1):70. <http://doi.org/10.1038/s41525-024-00451-7>
 27. Schnabel-Besson E, Mütze U, Dikow N, et al. Wilson and Jungner revisited: are screening criteria fit for the 21st century? *Int J Neonatal Screen.* 2024;10(3):62. <http://doi.org/10.3390/ijns10030062>
 28. Gordon AS, Lee K, Abul-Husn NS, et al. Consideration of disease penetrance in the selection of secondary findings gene-disease pairs: a policy statement of the American College of Medical Genetics and Genomics (ACMG). *Genet Med.* 2024;26(7):101142. <http://doi.org/10.1016/j.gim.2024.101142>
 29. Downie L, Bouffler SE, Amor DJ, et al. Gene selection for genomic newborn screening: moving toward consensus? *Genet Med.* 2024;26(5):101077. <http://doi.org/10.1016/j.gim.2024.101077>
 30. Minten T, Bick S, Adelson S, et al. Data-driven consideration of genetic disorders for global genomic newborn screening programs. *Genet Med.* 2025;27(7):101443. <http://doi.org/10.1016/j.gim.2025.101443>
 31. Betzler IR, Hempel M, Mütze U, et al. Comparative analysis of gene and disease selection in genomic newborn screening studies. *J Inherit Metab Dis.* 2024;47(5):945-970. <http://doi.org/10.1002/jimd.12750>
 32. Shang Z. Use of Delphi in health sciences research: a narrative review. *Medicine (Baltimore).* 2023;102(7):e32829. <http://doi.org/10.1097/MD.00000000000032829>
 33. Jünger S, Payne SA, Brine J, Radbruch L, Brearley SG. Guidance on Conducting and REporting DELphi Studies (CREDES) in palliative care: recommendations based on a methodological systematic review. *Palliat Med.* 2017;31(8):684-706. <http://doi.org/10.1177/0269216317690685>
 34. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research Electronic Data Capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42(2):377-381. <http://doi.org/10.1016/j.jbi.2008.08.010>
 35. Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform.* 2019;95:103208. <http://doi.org/10.1016/j.jbi.2019.103208>
 36. von der Gracht HA. Consensus measurement in Delphi studies: review and implications for future quality assurance. *Technol Forecast Soc Change.* 2012;79(8):1525-1536. <http://doi.org/10.1016/j.techfore.2012.04.013>
 37. Green RC, Shah N, Genetti CA, et al. Actionability of unanticipated monogenic disease risks in newborn genomic screening: findings from the BabySeq Project. *Am J Hum Genet.* 2023;110(7):1034-1045. <http://doi.org/10.1016/j.ajhg.2023.05.007>
 38. Pereira S, Smith HS, Frankel LA, et al. Psychosocial effect of newborn genomic sequencing on families in the BabySeq project: a randomized clinical trial. *JAMA Pediatr.* 2021;175(11):1132-1141. <http://doi.org/10.1001/jamapediatrics.2021.2829>
 39. Veldman A, Kiewiet MBG, Westra D, et al. A Delphi survey study to formulate statements on the treatability of inherited metabolic disorders to decide on eligibility for newborn screening. *Int J Neonatal Screen.* 2023;9(4):56. <http://doi.org/10.3390/ijns9040056>
 40. Kashtan CE, Gross O. Clinical practice recommendations for the diagnosis and management of Alport syndrome in children, adolescents, and young adults—an update for 2020. *Pediatr Nephrol.* 2021;36(3):711-719. <http://doi.org/10.1007/s00467-020-04819-6>
 41. Forrest IS, Chaudhary K, Vy HMT, et al. Population-based penetrance of deleterious clinical variants. *JAMA.* 2022;327(4):350-359. <http://doi.org/10.1001/jama.2021.23686>
 42. Gold JI, Madhavan S, Park J, et al. Phenotypes of undiagnosed adults with actionable OTC and GLA variants. *HGG Adv.* 2023;4(4):100226. <http://doi.org/10.1016/j.xhgg.2023.100226>
 43. Kingsmore SF, Wright M, Smith LD, et al. Prequalification of genome-based newborn screening for severe childhood genetic diseases through federated training based on purifying hyperselection. *Am J Hum Genet.* 2024;111(12):2618-2642. <http://doi.org/10.1016/j.ajhg.2024.10.021>